

Lemniscate of Bernoulli

Arseniy V. Akopyan

Abstract

In this paper we give purely geometrical proofs of the well-known properties of the lemniscate of Bernoulli.

What is lemniscate? *Polynomial lemniscate with foci $F_1, F_2, \dots, F_n$ is a locus of the points X with prescribed constant value of the product of distances from X to the foci. A root of degree n from this value is called a *radius* of lemniscate. Lemniscate is algebraic curve of degree (at most) $2n$. You can see the family of lemniscate with three foci on the figure 1.*




Fig. 1

Exercise 1. Write down a polynomial equation of lemniscate with n foci.

Lemniscate with two foci is called *Cassini oval*. The most interesting Cassini oval is a *Lemniscate of Bernoulli*, such curve that for every point on it product of distances from the foci equal quarter of square of distance between foci (fig. 2).

It's obvious that the lemniscate passes through the midpoint between foci. This point is called a *juncture* or *double point* of the lemniscate.




Fig. 2 $|F_1X| \cdot |F_2X| = |F_1O| \cdot |F_2O|$.

Lemniscate of Bernoulli has many very interesting properties. For example, area bounded by the lemniscate is equal to $\frac{1}{2}|F_1F_2|^2$. In this paper we prove some another properties, mainly using purely synthetic arguments.

How to construct Bernoulli's lemniscate? There exists a very simple method to construct Bernoulli's lemniscate using the following construction with three linked sticks. Suppose that two sticks F_1A and F_2B are linked in points F_1 and F_2 respectively, and have lengths equal to $\frac{1}{\sqrt{2}}|F_1F_2|$. Then let the points A and B lie on the different sides of the line F_1F_2 . The third sticks joins the points A and B and its length equals to $|F_1F_2|$ (fig. 3).




Fig. 3

Let us show, that under motion of this construction *midpoint of AB is moving on the lemniscate of Bernoulli with foci F_1 and F_2* .

Proof. Denote midpoint of the segment AB by X . Note, that polygon F_1AF_2B is a isosceles trapezoid (fig. 4). Moreover, triangles $\triangle AF_1X$ and $\triangle ABF_1$ are similar, because they have a common angle A and the following equation holds

$$\frac{|AF_1|}{|AX|} = \frac{|AB|}{|AF_1|} = \sqrt{2}.$$

For the same reason triangles $\triangle BXF_2$ and $\triangle BF_2A$ are similar. They have a common angle B , and the ratios of the length of the sides with endpoints in B equals to $\sqrt{2}$. Therefore, we can write the following equations:

$$\angle AF_1X = \angle ABF_1 = \angle BAF_2 = \angle XF_2B.$$




Fig. 4

Let us remark that in the trapezoid F_1AF_2B angles $\angle A$ and $\angle F_2$ are equal. Since angles $\angle XAF_2$ and $\angle XF_2B$ are equal too, we obtain that $\angle F_1AX = \angle XF_2A$. This implies that triangles $\triangle F_1AX$ and $\triangle AF_2X$ are similar. Therefore,

$$\frac{|F_1X|}{|AX|} = \frac{|AX|}{|XF_2|} \Rightarrow |XF_1| \cdot |XF_2| = |AX|^2 = |F_1O|^2.$$

It follows that the point X lies on the Bernoulli's lemniscate.

Note also that since the motion of the point X is continuous and X attains most distant points of the lemniscate, that trajectory of the point X is the whole lemniscate of Bernoulli.

Exercise 2. Suppose points A and B lie on the same side of the line F_1F_2 . Find the locus of the point X in this case.




Fig. 5

Suppose O is a midpoint of the segment F_1F_2 (double point of the lemniscate). Denote by M and N the midpoints of the segments F_1A and F_1B respectively (fig. 5). Shift the point O to the vector $\overrightarrow{NF_1}$. Denote obtained point by O' . Note that triangles $\triangle F_1MO'$ and $\triangle NXO$ are equal, moreover the following equation holds:

$$|F_1M| = |F_1O'| = \frac{1}{\sqrt{2}}|F_1O|.$$

In other words the points M and O' lie on the circle with center at F_1 and radius $\frac{1}{\sqrt{2}}|F_1O|$.

Using this we can obtain another elegant method to construct the Bernoulli's lemniscate.




Fig. 6

Let us construct the circle with center at one of the foci and radius $\frac{1}{\sqrt{2}}|F_1O|$. On each secant OAB (where A and B are the points of intersection of the circle and the secant) chose points X and X' , such that $|AB| = |OX| = |OX'|$ (fig. 6). Union of all points X and X' form a lemniscate of Bernoulli with foci F_1 and F_2 .




Fig. 7

Another interesting way to construct lemniscate with linkages you can see on the figure 7. Lengths of the segments F_1A and F_1O are equal to each other. The point A is the vertex of sticks AX and AY , and length of each of these sticks equals to $\sqrt{2}|F_1O|$. Denote the midpoints of these sticks by B and C , and join them with O by the sticks with length $\frac{|AX|}{2}$. In the process of rotating of point A around the circle each of the points X and Y generates a half of Bernoulli's lemniscate with foci F_1 and F_2 .

Exercise 3. Prove this.

Bernoulli's lemniscate and equilateral hyperbola. Hyperbola is a much more well-known curve. Hyperbola is a set of all points X such that value $||F_1X| - |F_2X||$ is a constant. Points F_1 and F_2 are called the foci of the hyperbola. Among all hyperbolas we single out a *equilateral* or *rectangular hyperbolas*. This is the set of points X such that $||F_1X| - |F_2X|| = \frac{|F_1F_2|}{\sqrt{2}}$.

Exercise 4. Show that $y = \frac{1}{x}$ is equation of equilateral hyperbolas and find the foci of it.

It appears that the lemniscate of Bernoulli is an inversion image of equilateral hyperbola. Before to prove this let us recall definition of inversion.

Definition 1. Inversion in the circle with center O and radius r is a transformation for which every point X in the plane maps to the point X^* lying on the ray OX and such that $|OX^*| = \frac{r^2}{|OX|}$.

Inversion has one very useful property: image of circles and lines are circles or lines again. About this and other very interesting properties of inversion you can read in [1]. We will prove here just one simple lemma that will be very helpful later.

Lemma 1. Suppose A is an orthogonal projection of the point O on some line l . Then inversion image of the line l in a circle with center O is a circle with diameter OA^* , where A^* is an inversion image of the point A .




Fig. 8

Proof. Let B be any point on the line l , and B^* its inversion image (fig. 8). Since

$$|OA^*| = \frac{r^2}{|OA|} \text{ and } |OB^*| = \frac{r^2}{|OB|},$$

we obtain that triangles $\triangle OAB$ and $\triangle OB^*A^*$ are similar. Therefore angle $\angle OB^*A^*$ is right and point B^* lies on the circle with diameter OA^* . $\square$

It is necessary to note that center O_1^* of this circle is the inversion image of the point O_1 , where O_1 is the image of point O after symmetry in line l .

Now, let us prove that *lemniscate of Bernoulli with foci F_1 and F_2 is an inversion image of equilateral hyperbola with foci F_1 and F_2 with respect to the circle with center O and radius $|OF_1|$*

For this we will use the results that we obtain in the proof of robustness of the first method to construct the lemniscate (fig. 4). Denote by P point of intersection of the lines F_1A and F_2B and let Q be its image after symmetry in the line F_1F_2 (fig. 9). Let us note that

$$|F_2Q| - |F_1Q| = |F_2P| - |F_1P| = |AP| - |F_1P| = |F_1A| = \frac{|F_1F_2|}{\sqrt{2}}.$$

Therefore, points P and Q lie on the equilateral hyperbola with foci at F_1 and F_2 . Now is left to show that point X and Q is inversion image of each other with respect to the circle with center O and radius $|OF_1|$. First, let us show that triangles $\triangle F_1XO$ and $\triangle PF_1O$ are similar.




Fig. 9

Note that F_1XOB is a trapezoid, therefore $\angle OXF_1 + \angle XF_1B = 180^\circ$. Also we have $\angle AF_1O + \angle OF_1P = 180^\circ$. Since angle $\angle XF_1B$ is equal to angle $\angle AF_1O$, we obtain that $\angle OXF_1$ and $\angle OF_1P$ are equal to each other.

Since angles $\angle XF_2B$ and $\angle XF_1A$ are equal, we have that $\angle XF_1P + \angle PF_2X = 180^\circ$, in other words quadrilateral PF_1XF_2 is inscribed. Therefore

$$\angle F_2F_1X = \angle F_2PX = \angle F_1PO.$$

The last equation holds as far as point O and X are symmetry to each other with respect to perpendicular bisector to the segment F_1B .

Thus, triangles $\triangle F_1XO$ and $\triangle PF_1O$ are similar because have two corresponding pairs of equals angles. It follows that $\angle F_1OX$ and $\angle F_1OP$ are equal, and we have that the point Q lies on the ray OX . As well, from similarity of triangles $\triangle F_1XO$ and $\triangle QF_1O$ (it is congruent to the triangle $\triangle PF_1O$) we obtain that

$$\frac{|OX|}{|OF_1|} = \frac{|OF_1|}{|OQ|} \Rightarrow |OX| \cdot |OQ| = |OF_1|^2.$$

It means that points Q and X are inversion images of each other with respect to the circle with center O and radius $|OF_1|$.

If we look on the figure 9 we can make another observation: the points X and O lie on the circle with center P . It is interesting that *this circle touches¹ the lemniscate of Bernoulli.*




Fig. 10

Proof. Suppose l is a tangent line to the hyperbola in the point Q . From Lemma 1 it follows that image of the line l after inversion with respect to the circle with center O and radius $|F_1O|$ is a circle ω_l which passes through the point O . Since X is an inversion image of the point Q , we see that this circle ω_l touches the lemniscate at the point X . From the same Lemma we can conclude that center of this circle lies on the normal line from the point O to the line l . Let us show that lines OP and OQ is symmetric to each other with respect to the line F_1F_2 . It will follow that the point P is a center of the circle ω_l .




Fig. 11

Without loss of generality we can assume that equation of the hyperbola is $y = \frac{1}{x}$. Suppose line l intersects the abscissa and the ordinate in the points R and S respectively (fig. 11). It is well-known that derivative of function $\frac{1}{x}$ in the point x_0 is equal to $-\frac{1}{x_0^2}$. It follows that the point Q is a midpoint of the segment RS , and OQ is a median of the right triangle $\triangle ROS$, therefore angles $\angle QOR$ and $\angle QRO$ are equal. Since angles $\angle POS$ and $\angle QOR$ are equal too, we obtain that the lines OP and RS are perpendicular, as was to be proved. $\square$




Fig. 12

¹Let us recall that two curves on the plane touch in a point if there exist line touching both curves in this point

Let us note the following. Since circle ω_l tangents to the lemniscate in the point X , the radius PX of this circle is a normal (perpendicular to the tangent line) to the lemniscate in the point X (fig. 12). Note that the triangle $\triangle XPO$ is isosceles, and lines XO and PO are symmetric with respect to the line F_1O . Therefore we can write these equations:

$$\angle PXO = \angle XOP = 2\angle POF_1.$$

It follows that there exists the following very simple method to construct the normal to the lemniscate of Bernoulli. Indeed for any point X on the lemniscate construct the lines which with the line OX form the angle equal to the $2\angle XOF_1$. This line will be a normal to the lemniscate.

References

- [1] R. A. Johnson. Advanced Euclidean Geometry, 1925, 2007.
- [2] J.D. Lawrence. *A catalog of special plane curves*. Dover Publications New York, N. Y, 1972.
- [3] A.I. Markushevich. *Zamechatelniye kriviye*. M.:Gostekhizdat, 1952.